



Bronllys Hospital, Bronllys, Brecon, Powys, LD3 0LU

This Patient Group Direction (PGD) must only be used by registered healthcare professionals who have been named and authorised by their organisation to practice under it. The most recent and in date final signed version of the PGD should be used. Healthcare professionals should always access the PGD via the PTHB internet to ensure that they are always working to the most up to date version.

Patient Group Direction

for the administration of

Measles, mumps, rubella and varicella (MMRV) vaccine

by

registered Healthcare Professionals

to

individuals from one year of age and eligible for routine or catch-up immunisation,

or

from 9 months of age if early protection is required against varicella

in

Powys Teaching Health Board

Version number: PGD 0250A

Change History

Version number	Change details	Date
PGD 0250	Adopted from the new UKHSA MMRV PGD v1.0 template to support the delivery of the national combined measles, mumps, rubella and varicella immunisation programme to eligible children between the ages of one and under 6 years of age.	18/11/2025
PGD 0250A	<p>Review issue in line with UKHSA MMRV PGD template v2.0:</p> <ul style="list-style-type: none"> • clarification added to inclusion criteria, cross-referencing the advice in dose and frequency of administration, that children who have missed vaccination for MMR born on or after 1 January 2020 are in scope of this PGD • recommendation that individuals born on or after 1 January 2020, who have previously received 2 valid doses of MMR vaccine and who are not captured in the selective MMRV catch-up programme, should receive a single dose of MMRV vaccine • off-label section advice updated regarding individuals taking regular salicylates who receive a live varicella-containing vaccine, as outlined in the live monovalent varicella vaccine PGD 0198 • clarification in the dose and frequency of administration section that children born on or after 1 January 2020 and on or before 31 August 2022, who have missed routine MMR doses must not be deferred until the start of the selective MMRV catch-up programme starting 1 November 2026; offer up to 2 MMRV doses • signposting to the MMRV eligibility tables as an additional resource for those with missing MMRV doses • clarification of interval between first and second MMR-containing dose (amended from one month to 4 weeks, in line with other MMRV programme documents) • clarification that an additional MMRV dose is required at 18 months if further MMR-containing doses were offered less than 3 months after the first dose and when aged under 18 months of age (previously 	02 /02/2026

Reference Number: PGD 0250A

Valid from: 02/02/2026

Review Date: 31/05/2028

Expiry Date: 30/11/2028

	<p>advised second doses offered before 15 months of age should be repeated)</p> <ul style="list-style-type: none"> • minor correction made to Appendix and the brand names of monovalent varicella vaccine (Varivax® and Varilrix®, not Vaxelis® and Varilrix® as previously published) <p>Reviewed to include minor rewording of standard text, layout and formatting changes for clarity and consistency with other PTHB PGDs. Title amended to match PGD template more closely.</p>	
--	---	--

his Powys Teaching Health Board (PTHB) PGD is based on the UKHSA MMRV vaccine PGD template v2.0 developed by the following health professionals on behalf of the UKHSA and peer reviewed by the UKHSA Immunisations PGD Expert Panel in accordance with the UKHSA PGD and Protocol Policy (also ratified by the UKHSA Medicines Governance Committee). The UKHSA template has been adapted for use in PTHB.





Developed by the following health professionals on behalf of the UKHSA:

Developed By:	Name
Pharmacist (Lead Author)	Christina Wilson, Lead Pharmacist - Immunisation Programmes, UKHSA
Doctor	Dr Vanessa Saliba Consultant Medical Epidemiologist, Immunisation and Vaccine Preventable Diseases Division, UKHSA
Registered Nurse (Chair of Expert Panel)	David Green Nurse Consultant Immunisation Programmes - UKHSA

Expert Panel

Name	Designation
Nicholas Aigbogun	Consultant in Communicable Disease Control, Yorkshire and Humber Health Protection Team, UKHSA
Jess Baldasera	Health Protection Practitioner, Northeast Health Protection Team, Regions Directorate, UKHSA
Helen Beynon	Clinical Advisor, Immunisation Clinical Advice Response Service (CARS), NHS England – London
Alison Campbell	Screening and Immunisation Coordinator, Clinical, NHS England – Midlands
Laura Craig	Lead Immunisation Nurse Specialist, Immunisation Programmes –UKHSA
Jane Freeguard	Deputy Director of Vaccination – Medicines and Pharmacy, NHSE
Rosie Furner	Advanced Specialist Pharmacist – Medicines Governance (Patient Group Directions and Medicines Mechanisms), NHS Specialist Pharmacist Services (SPS)
Ed Gardner	Advanced Paramedic Practitioner/ Emergency Care Practitioner, Primary Care Based, Southbourne Surgery
Shilan Ghafoor	Lead Medicines Governance Pharmacist, Medicines Governance, UKHSA
Michelle Jones	Principal Medicines Optimisation Pharmacist, NHS Bristol, North Somerset and South Gloucestershire Integrated Care Board
Elizabeth Lockett	Senior Screening and Immunisation Manager, Screening and Immunisation Team – Kent and Medway, NHS England – South East
Dr Vanessa MacGregor	Consultant in Communicable Disease Control, East Midlands Health Protection Team, UKHSA
Lesley McFarlane	Lead Immunisation Nurse Specialist, Immunisation Programmes –UKHSA
Briony Mason	Vaccination Manager and Professional Midwifery Advocate, Vaccination and Screening, NHS England – West Midlands
Tushar Shah	Lead Pharmacy Adviser, NHS England – London

PGD Authorisation

Name	Job title and organisation	Signature	Date
Senior doctor Dr Kate Wright	Lead doctor for PTHB	 DocuSigned by: Kate Wright 1F267952823F473...	2/23/2026
Chief Pharmacist Jonathan Boyd	Chief Pharmacist for PTHB	 Signed by: Jon Boyd 6D8ECFE8C9EB423...	2/10/2026
Senior representative of professional group using the PGD Paul Hooton	Executive Director of Nursing and Midwifery for PTHB	 Signed by: Paul Hooton EEABC83AC83F4B9...	2/9/2026
Clinical Governance Lead Amanda Edwards	Clinical Governance Lead for PTHB –Assistant Director for Innovation and Improvement	 DocuSigned by: Amanda Edwards 74A4E51A42E9473...	3/3/2026

The PGD is not legally valid until it has had the relevant organisational authorisation. [Appendix A](#) provides a table listing PGD indications for use of MMR, MMRV and monovalent varicella vaccine, in accordance with the individual's age.

It is the responsibility of the organisation that has legal authority to authorise the PGD, to ensure that all legal and governance requirements are met. The authorising body accepts governance responsibility for the appropriate use of the PGD.

[Appendix B](#) provides a staff accreditation sheet. Individual practitioners must be authorised by name, under the current version of this PGD, before working to it.

Those using this PGD must ensure that it is organisationally authorised and signed by an appropriate authorising person, relating to the class of person by whom the product is to be supplied, in accordance with Human Medicines Regulations 2012 (HMR2012)¹. **The PGD is not legal or valid without signed authorisation in accordance with [HMR2012 Schedule 16 Part 2](#).**

Operation of this PGD is the responsibility of commissioners and service providers. The final authorised copy of this PGD should be kept by PTHB for 25 years after the

¹ This includes any relevant amendments to legislation

Reference Number: PGD 0250A

Valid from: 02/02/2026

Review Date: 31/05/2028

Expiry Date: 30/11/2028

PGD expires. Provider organisations adopting authorised versions of this PGD should also retain copies for 25 years after the PGD expires.

Individual practitioners must be authorised by name, under the current version of this PGD before working according to it.

Practitioners and organisations must check that they are using the current version of the PGD. Amendments may become necessary prior to the published expiry date. Enquiries relating to the availability of organisationally authorised PGDs and subsequent versions of this PGD should be directed to:
info.medicinesmanagement.powys@wales.nhs.uk

PGD adoption by the provider

Name	Job title and organisation	Signature	Date

Reference Number: PGD 0250A
Valid from: 02/02/2026
Review Date: 31/05/2028
Expiry Date: 30/11/2028

1. Characteristics of staff

<p>Qualifications and professional registration</p>	<p>All practitioners should only administer vaccinations where it is within their clinical scope of practice to do so. Practitioners must also fulfil the additional requirements and continued training requirements to ensure their competency is up to date, as outlined in the sections below.</p> <p>Practitioners working to this PGD must also be one of the following currently registered professionals who can legally supply and administer under a PGD:</p> <ul style="list-style-type: none"> • nurses and midwives currently registered with the Nursing and Midwifery Council (NMC) • pharmacists and pharmacy technicians currently registered with the General Pharmaceutical Council (GPhC) (Note: this PGD is not relevant to privately provided community pharmacy services) • dieticians, occupational therapists, paramedics, physiotherapists and podiatrists currently registered with the Health and Care Professions Council (HCPC) <p>Check Appendix B – Staff Accredited to use Patient Group Direction to confirm whether all practitioners listed above have organisational authorisation to work under this PGD.</p>
<p>Additional requirements</p>	<p>Additionally, practitioners:</p> <ul style="list-style-type: none"> • must be authorised by name as an approved practitioner under the current terms of this PGD before working to it • must have undertaken appropriate training for working under PGDs for supply/administration of medicines. Must have completed eLfh PGD eLearning Patient Group Directions training (available via learning@wales, PTHB staff to access via ESR). Evidence of ongoing PGD training to be submitted to Line Manager annually– this should include an annual completion certificate of PGD e-learning or a dated screenshot of the PGD e-learning assessment results as proof of completion. • must be competent in the use of PGDs (see NICE Competency framework for health professionals using PGDs). Individuals operating under this PGD must be assessed as competent (see Appendix B) • must be familiar with the vaccine product and alert to changes in the Summary of Product Characteristics (SPC), Immunisation Against

	<p>Infectious Disease (the Green Book) and national and local immunisation programmes</p> <ul style="list-style-type: none"> • must have undertaken training appropriate to this PGD as required by local policy and in line with the National Minimum Standards and Core Curriculum for Immunisation Training and online training. Please contact PTHB immunisation co-ordinator for further information. • must be competent to undertake immunisation and to discuss issues related to immunisation • must be competent in the handling and storage of vaccines, and management of the cold chain. Completion of cold chain training (also available via ESR) • must be familiar with All Wales Advisory document on Ordering Storage and Handling of Vaccines • must be competent in the appropriate administration method for the vaccines listed in this PGD • must be competent in the recognition, management and reporting of adverse drug reactions, including anaphylaxis. Must be competent in the administration of adrenaline 1 in 1000 and have up to date Basic Life Support (BLS) skills • must have access to the PGD and associated online resources • should fulfil any additional requirements defined by local policy <p>THE INDIVIDUAL PRACTITIONER MUST BE AUTHORISED BY NAME, UNDER THE CURRENT VERSION OF THIS PGD BEFORE WORKING ACCORDING TO IT.</p>
<p>Continued training requirements</p>	<p>Practitioners must ensure they are up to date with relevant issues and clinical skills relating to immunisation and management of anaphylaxis, with evidence of appropriate Continued Professional Development (CPD) (updating at least every 2 years on the administration of MMRV vaccine).</p> <p>Practitioners must be competent, recognise their own limitations and personal accountability and act accordingly. Practitioners must make a self-declaration of competency on PADR (if relevant). The personal development plan (yellow) section of the PADR booklet should be used to record completion of Statutory and Mandatory training, including annual PGD e-learning.</p>

	<p>Compliance with all mandatory NHS training (if relevant).</p> <p>Practitioners should be constantly alert to any subsequent recommendations from the Welsh Government and/or Public Health Wales and/or NHS Wales and/or the UKHSA and/or NHS England and/or other sources of medicines information.</p> <p>Note: the most current national recommendations should be followed but a Patient Specific Direction (PSD) may be required to administer the vaccine in line with updated recommendations that are outside the criteria specified in this PGD.</p> <p>It is the responsibility of the healthcare professional to maintain their own competency to practice within this PGD.</p>
--	--

2. Clinical condition or situation to which this PGD applies

<p>Clinical condition or situation to which this PGD applies</p>	<p>Indicated for the active immunisation of individuals from one year of age for routine or catch-up immunisation, or from 9 months of age if early protection is required against varicella and recommendations given in the measles, mumps, rubella and varicella chapters of Immunisation Against Infectious Disease: the Green Book.</p> <p>Note: appendix A provides an overview of the scope of this and the MMR PGD0163.</p> <p>If used in PTHB Occupational Health Service for their own staff (including students and trainees), the PGD must be used in line with PTHB Occupational Immunisation policy.</p> <p>It is the responsibility of the administering healthcare professional to ensure that the individual is within the inclusion criteria, and that there are no reasons for exclusion before proceeding with immunisation. If there is any reason for concern, seek medical advice.</p>
---	--

Criteria for inclusion	<p>1. Routine MMRV programme</p> <ul style="list-style-type: none"> all children presenting for a measles-containing vaccine with a date of birth (DOB) on or after 1 January 2020 <p>This includes:</p> <ul style="list-style-type: none"> children who are late, have missed scheduled doses or who are unvaccinated, incompletely vaccinated or have an unknown MMR vaccination status <p>Vaccination should not be delayed for those who have missed scheduled doses.</p> <p>2. Selective MMRV catch-up programme</p> <p>A one-dose MMRV selective catch-up programme will be delivered between 1 November 2026 to 31 March 2028 (or other end date as subsequently communicated by NHS England/Wales).</p> <ul style="list-style-type: none"> children aged from 3 years 4 months to under 6 years on 31 December 2025 (DOB on or after 1 January 2020 to 31 August 2022) without a history of chickenpox disease or 2 doses of varicella vaccine are eligible for one dose of MMRV <p>3. children born on or after 1 January 2020 who have received 2 valid doses of MMR vaccine and who are not included in criterion 2 should be offered a single dose of MMRV vaccine.</p> <p>To provide urgent measles or varicella protection, or opportunistic MMR catch up</p> <p>4. when the child is in one of the eligible cohorts as outlined above (aged 12 months and over) and protection is indicated as part of a response to a measles outbreak or for travel to a measles endemic area.</p> <p>5. when an individual aged 9 months of age and over requires urgent protection against either varicella or measles, such as during an outbreak and is eligible for vaccination with the varicella vaccine (refer to the varicella vaccine PGD 0198 Patient Group Directions (PGDs) - Powys Teaching Health Board) or for vaccination with MMR (refer to the MMR PGD 0163 Patient Group Directions (PGDs) - Powys Teaching Health Board) but no stock of either is available.</p>
-------------------------------	--

Reference Number: PGD 0250A

Valid from: 02/02/2026

Review Date: 31/05/2028

Expiry Date: 30/11/2028

	<p>6. when an individual aged 9 months of age and over requires both MMR and monovalent varicella vaccination at the same time, it is more pragmatic to give the MMRV vaccine.</p> <p>7. when an individual not eligible for MMRV requires an opportunistic dose of MMR vaccine to bring their vaccination history up to date, but there is no stock of the MMR vaccine available.</p> <p>Informed consent, from the individual or a person legally able to act on the individual’s behalf, must be obtained prior to administration. NB. Refer to PTHB Consent to Treatment and Examination Policy.</p> <p>Medical and drug history taken, no reason for exclusion.</p> <p>Any vulnerable adult or child protection concerns should be referred to Safeguarding and PTHB safeguarding policies followed. Where there are safeguarding concerns (Child Protection or Protection of Vulnerable Adults, POVA) advice from the local Safeguarding team should be sought (see below).</p>
<p>Criteria for exclusion Exclusion under the PGD does not necessarily mean the medication is contraindicated, but it would be outside its remit and another form of authorisation will be required</p>	<p>Individuals for whom no valid consent has been received (or for whom a best-interests decision in accordance with the Mental Capacity Act 2005, has not been obtained). For further information on consent, see chapter 2 of the Green Book. Several resources are available to inform consent (see written information to be given to individual, parent or carer section). If consent is not received, please refer to section Action to be taken if the individual, parent or carer declines treatment.</p> <p>Individuals who:</p> <ul style="list-style-type: none"> • have had a confirmed anaphylactic reaction to a previous dose of any measles, mumps, rubella or varicella containing vaccine or to any components of the vaccine. These include neomycin and sorbitol (refer to relevant SPCs) • were born on or before 31 December 2019, except those who require vaccination under points 4, 5, 6 or 7 of the criteria for inclusion above • are under 9 months of age • are known to be pregnant

	<ul style="list-style-type: none"> • have a primary or acquired immunodeficiency state (see the Green Book chapter 6 for more detail) • are on current or recent high dose immunosuppressive or biological therapy (see the Green Book chapter 6 for more detail) • have received a live varicella-containing vaccine or yellow fever vaccine in the preceding 4 weeks, unless protection against measles is rapidly required (see drug interactions) • have received blood products, such as immunoglobulins, in the preceding 3 months, unless protection against measles is rapidly required (see drug interactions) • are awaiting reading of a tuberculin (Mantoux) skin test, unless protection against measles is rapidly required (see drug interactions) • are suffering from acute severe febrile illness (the presence of a minor infection is not a contraindication for immunisation) • have received 2 doses of MMRV, at an appropriate age to be effective (see also special considerations and additional information – early vaccination due to travel, outbreak or contact with a probable or confirmed case of measles) • Conditions outside of the clinical situations criteria. <p>Please refer to section Action to be taken if the individual is excluded.</p>
<p>Cautions including any relevant action to be taken</p>	<p>Facilities for management of anaphylaxis should be available at all vaccination sites (see Chapter 8 of the Green Book) and advice issued by the Resuscitation Council UK.</p> <p>Individuals who are immunosuppressed or who are living with HIV, who are not contraindicated to receive this live vaccine (see the Green Book chapter 6) may not make a full antibody response and revaccination upon cessation of treatment or clinical recovery may be required. This should be discussed with the appropriate specialist and the repeat dose administered under a PSD.</p> <p>If idiopathic thrombocytopenic purpura (ITP) has occurred within 6 weeks of the first dose of MMRV, then blood should be taken and tested for measles antibodies before a second dose is given. Serum should be sent to the UKHSA Virus Reference</p>

Department, which offers free, specialised serological testing for such children. If the results suggest incomplete immunity against measles, then a second dose of MMRV is recommended.

The presence of a neurological condition is not a contraindication to immunisation but if there is evidence of current neurological deterioration, deferral of vaccination may be considered, to avoid incorrect attribution of any change in the underlying condition. The risk of such deferral should be balanced against the risk of the preventable infection, and vaccination should be promptly given once the diagnosis or the expected course of the condition (or both) become clear. There will be very few occasions when deferral of immunisation is required. Deferral leaves the child unprotected and so the period of deferral should be minimised, with immunisation commencing as soon as possible. If a specialist recommends deferral, this should be clearly communicated to the individual's primary care provider, who must be informed as soon as the child is fit for immunisation. Children with a personal or close family history of seizures should still be given the MMRV vaccine.

Priorix Tetra[®] and ProQuad[®] both contain a source of phenylalanine. Though phenylalanine may be harmful to individuals with phenylketonuria (PKU), such individuals (or their parent or carer) will be well versed as to the amounts of phenylalanine tolerable in their diet. The National Society for Phenylketonuria (NSPKU) advise the amount of phenylalanine contained in vaccines is negligible and therefore strongly advise individuals with PKU to take up the offer of immunisation.

Syncope (fainting) can occur following, or even before, any vaccination especially in adolescents as a psychogenic response to the needle injection. This can be accompanied by several neurological signs such as transient visual disturbance, paraesthesia and tonic-clonic limb movements during recovery. It is important that procedures are in place to avoid injury from faints.

Check for any other medications that the individual is taking, including topical or inhaled products, food supplements and herbal or homeopathic products. (Refer to [BNF/SPC](#) for full list).

Reference Number: PGD 0250A

Valid from: 02/02/2026

Review Date: 31/05/2028

Expiry Date: 30/11/2028

	<p>Under Section 128 and 130 of the Social Services and Wellbeing (Wales) Act 2014, staff have a duty to inform the Local Authority if they have reasonable cause to suspect that an adult or child is at risk. Any vulnerable adult or child protection concerns should be referred to Safeguarding and the PTHB safeguarding policies followed. Consider discussing with GP. Any safeguarding concerns need to be directed to Safeguarding Hub:</p> <p>To generic email address: PowysTHB.Safeguarding@wales.nhs.uk and</p> <ul style="list-style-type: none"> • Central Safeguarding number: 01686 252806 • Out of hours: 0345 0544847 <p>Advice can also be sought from local Safeguarding Leads</p>
<p>Action to be taken if the individual is excluded</p>	<p>Explain reason to individual, parent or carer.</p> <p>Individuals who have had a confirmed anaphylactic reaction to a previous dose of MMRV vaccine or to any components of the vaccine should be referred to a clinician for specialist advice and appropriate management.</p> <p>Individuals who are pregnant should be advised to avoid contact with known or suspected cases of measles, mumps, rubella or varicella infection and report any rash illness or contact with rash illness to their GP or midwife (or both). Women who are lacking 2 documented doses of MMR should be immunised after their pregnancy, at the earliest opportunity and before any further pregnancies. Note: MMRV can be given to breast-feeding mothers without any risk to their baby.</p> <p>Individuals who have a primary or acquired immunodeficiency state or who are currently, or were recently, on high dose immunosuppressive or biological therapy (see chapter 6) should consult the appropriate specialist regarding the individual’s immune status and suitability for receiving live MMRV vaccine. Where administration of MMRV is advised, a PSD will be required. Further information to guide suitability of the MMRV vaccine for individuals living with HIV is available in Table 1 (CD4 count), of the Green Book chapter on measles.</p>

Reference Number: PGD 0250A
Valid from: 02/02/2026
Review Date: 31/05/2028
Expiry Date: 30/11/2028

Individuals who have been immunised against varicella or yellow fever within the last 4 weeks, or received blood products in the preceding 3 months, and do not require rapid protection against measles, mumps or rubella, should defer immunisation until the appropriate minimum interval has been observed (see [drug interactions](#) section).

Individuals who are awaiting reading of a tuberculin (Mantoux) test should delay vaccination with MMRV until the skin test has been read, unless protection against measles is required urgently.

Individuals aged 6 years and over on 1 January 2026, with a DOB on or before 31 December 2019 are not routinely in scope for inclusion with the MMRV programme. If the individual has not had 2 doses of MMR vaccine, they should complete the course with the MMR vaccine. Refer to the [MMR PGD0163](#). The main exception to this is in cases (as per [criteria for inclusion](#)) where there is no stock of MMR or of the monovalent varicella vaccine accessible and urgent protection is required, such as in post-exposure measles prophylaxis or where an opportunistic dose of MMR is required for an unimmunised or partially immunised individual. In such exceptions, MMRV may be given under this PGD.

The other main exception is in the rare event that an individual is identified as requiring both varicella and MMR vaccination. It is considered pragmatic to offer MMRV in such circumstances.

In case of postponement due to acute severe febrile illness, advise when the individual can be vaccinated and ensure another appointment is arranged.

Seek appropriate advice from the local Screening and Immunisation Team, local Health Protection Team or the individual's clinician as appropriate.

The risk to the individual of not being immunised must be taken into account.

Document the reason for exclusion and any action taken in the individual's clinical records.

Inform or refer to the GP or a prescriber as appropriate.

Reference Number: PGD 0250A

Valid from: 02/02/2026

Review Date: 31/05/2028

Expiry Date: 30/11/2028

	For PTHB Occupational Health Service- refer to Occupational Health Consultant as necessary and document advice given
Action to be taken if the individual, parent or carer declines treatment	<p>Advise the individual, parent or carer about the protective effects of the vaccine, the risks of infection and potential complications.</p> <p>Document the advice given and the decision reached.</p> <p>Inform or refer to the GP or a prescriber as appropriate.</p> <p>Inform child health if appropriate – if any vaccination is declined for a child under 19 years, child health must be informed, and the appropriate form completed. Where appropriate, inform the GP using the local agreed system.</p> <p>For PTHB Occupational Health Service- refer to Occupational Health Consultant as necessary and document advice given.</p>
Arrangements for referral for medical advice	<p>Refer to GP, paediatrician, consultant in communicable disease control (CCDC) or Occupational Health Consultant for clinical advice as necessary.</p> <p>Document any advice given.</p>

3. Description of treatment

Name, strength and formulation of drug	<p>Measles, mumps, rubella and varicella vaccine (live):</p> <ul style="list-style-type: none"> • ProQuad® powder and solvent for suspension for injection • Priorix Tetra® powder and solvent for solution for injection
Legal category	Prescription only medicine (POM)
Black triangle▼	No.
Off-label use	The national MMRV programme is scheduled at 12 and 18 months of age. This interval of 6 months is off label, as the SPCs recommend a minimum interval of one month (6 weeks for Priorix Tetra®) and preferably 3 months between doses, but represents national guidance.

Reference Number: PGD 0250A

Valid from: 02/02/2026

Review Date: 31/05/2028

Expiry Date: 30/11/2028

	<p>The SPC for Priorix Tetra® recommends that salicylates are avoided for 6 weeks following varicella vaccination. Individuals may continue with their salicylate treatment before and after varicella vaccination in accordance with varicella chapter of the Green Book.</p> <p>The vaccines should be stored according to the conditions detailed in the storage section below. However, in the event of an inadvertent or unavoidable deviation of these conditions, refer to the All Wales Advisory document on Ordering Storage and Handling of Vaccines and Vaccine Incident Guidance. Where vaccines are assessed in accordance with these guidelines as appropriate for continued use, this would constitute off-label administration under this PGD.</p> <p>Where a vaccine is recommended off-label consider, as part of the consent process, informing the individual, parent or carer that the vaccine is being offered in accordance with national guidance but outside of product licence.</p>
<p>Route and method of administration</p>	<p>The vaccine must be reconstituted in accordance with the manufacturer’s instructions prior to administration.</p> <p>Administer by intramuscular injection, preferably into the anterolateral aspect of the thigh in infants under one year of age. The deltoid muscle of the upper arm may be used in individuals over one year of age. (see The Green Book Chapter 4).</p> <p>When administering at the same time as other vaccines, care should be taken to ensure that the appropriate route of injection is used for all the vaccinations. The vaccines should be given at separate sites, preferably into different limbs. If given into the same limb, they should be given at least 2.5cm apart. The site at which each vaccine was given should be noted in the individual’s records.</p> <p>Individuals with bleeding disorders may be vaccinated intramuscularly if, in the opinion of a clinician familiar with the individual’s bleeding risk, vaccines or similar small volume intramuscular injections can be administered with reasonable safety by this route. Individuals on stable anticoagulation therapy, including individuals on warfarin who are up to date with their scheduled INR testing and whose latest INR was below</p>

Reference Number: PGD 0250A
 Valid from: 02/02/2026
 Review Date: 31/05/2028
 Expiry Date: 30/11/2028

	<p>the upper threshold of their therapeutic range, can be vaccinated via the intramuscular route. If the individual receives medication or other treatment to reduce bleeding, for example treatment for haemophilia, intramuscular vaccination can be scheduled shortly after such medication or other treatment is administered. A fine needle (equal to 23 gauge or finer calibre such as 25 gauge) should be used for the vaccination, followed by firm pressure applied to the site (without rubbing) for at least 2 minutes. The individual, parent or carer should be informed about the risk of haematoma from the injection.</p> <p>Both MMRV vaccines are licensed to be given subcutaneously, therefore a healthcare professional may determine this is a preferred route of administration for an individual with a bleeding disorder. Note fewer injection site reactions were reported with the intramuscular route compared with the subcutaneous route following administration of ProQuad®.</p> <p>The vaccine should be visually inspected for foreign particulate matter and other variation of expected appearance prior to preparation and administration. Should either occur, do not administer the dose and discard the vaccine in accordance with local procedures.</p> <p>Upon reconstitution, ProQuad® forms a clear pale yellow to light pink liquid. Priorix® Tetra forms a clear peach to fuchsia pink solution.</p> <p>The vaccine SPC provides further guidance on preparation and administration.</p> <p>Check expiry date and that the correct product has been chosen.</p>
<p>Dose and frequency of administration</p>	<p>Single 0.5ml dose per administration.</p> <p>The dosing schedules below assume individuals have been immunised in line with the complete routine immunisation schedule, without undue delay.</p> <p>For individuals who have missed routine appointments, please refer to information for healthcare professionals - Appendix A: MMRV eligibility tables (on the on gov.uk website), the PHW MMRV Eligibility Tool, and</p>

Reference Number: PGD 0250A

Valid from: 02/02/2026

Review Date: 31/05/2028

Expiry Date: 30/11/2028

[vaccination of individuals with uncertain or incomplete immunisation status](#) guidance and offer up to 2 MMRV doses accordingly.

Table 1: Routine MMRV programme and number of doses

	One dose schedule	2 dose schedule
DOB on or after 1 January 2025		12 months and 18 months
DOB on or after 1 July 2024 to 31 December 2024		18 months and 3 years 4 months
DOB on or after 1 September 2022 to 30 June 2024	3 years 4 months	

Table 2: Selective catch-up programme (1 November 2026 to 31 March 2028)

	One dose of MMRV*
DOB 1 January 2020 to 31 August 2022	Children without a history of chickenpox disease or 2 doses of varicella vaccination. There is no requirement to check the clinical history.

***Note:** those who have had 2 doses of the varicella vaccine privately may still receive the catch-up dose.

MMRV is not required if a child has had 2 MMR vaccines and either a history of previous chickenpox infection or 2 doses of varicella vaccine. The MMRV dose may still be given even if the child has had previous chickenpox infection or 2 doses of varicella vaccine. See also [special considerations and additional information – individuals who have already received live \(monovalent\) varicella vaccine \(including private market vaccination\)](#).

Incomplete immunisation history

Those born on or after 1 January 2020 overdue any doses of MMR-containing vaccine should be given up to 2 MMRV doses to ensure full protection against measles, mumps and rubella. The individual should be brought up to date at the earliest opportunity if routine appointments have been missed.

Please also refer to MMRV [eligibility tables](#) and [vaccination of individuals with uncertain or incomplete immunisation status](#) guidance.

In exceptional circumstances, there may be children with a DOB from 1 January 2020 not already included in the above schedules, who have already received 2 valid doses of MMR vaccine, but who are unimmunised against varicella. For example, there may be younger children who were vaccinated early due to travel, an outbreak or where vaccination was routinely offered at 18 months, not 3 years 4 months. In such cases, children should be offered a single dose of MMRV.

Incomplete immunisation history: children with a DOB between 1 January 2020 and 31 August 2022

Children who have missed doses of MMR-containing vaccine with a DOB from 1 January 2020 to 31 August 2022, should not have vaccination deferred until the start of the selective MMRV catch-up programme.

These children should be brought up to date at the earliest opportunity, with a minimum interval of 4 weeks between MMRV doses where applicable.

Children who have had no previous doses of MMR-containing vaccine, who receive 2 MMRV doses in this way do not need to be recalled for the selective MMRV catch-up programme.

There are no safety concerns if a child is given a total of 3 MMR-containing vaccines.

Early vaccination due to travel, outbreak or contact with a probable or confirmed case of measles or varicella

The MMRV vaccine can be given from 9 months of age when early protection is required against varicella. MMRV should not be given to individuals below 9 months of age. Vaccination for travel or measles outbreaks in this age group should be with MMR (see also [appendix A](#)).

- doses of MMRV given before 12 months of age should be discounted.
- additional doses given before 18 months of age should be given at least 3 months apart (providing the first dose was given from 12 months of age) and should be repeated from 18 months if this is not possible
- doses given from 18 months of age may be given at a 4 week interval

See [special considerations and additional information \(early vaccination due to travel, outbreak or contact](#)

	<p>with a probable or confirmed case of measles) for further information.</p> <p>In cases of post-exposure measles or varicella vaccination, the dose should ideally be given within 3 days of exposure to maximise vaccine efficacy.</p>
Duration of treatment	<p>Up to 2 doses of 0.5ml at the recommended interval (see dose and frequency of administration above).</p> <p>Doses that are administered early (see above) or doses given within 4 weeks of previous yellow fever, live varicella-containing vaccine or within 3 months of receiving blood products may need to be repeated (see drug interactions section and dose and frequency of administration above).</p>
Quantity to be administered	<p>Single 0.5ml dose per administration.</p>
Supplies	<p>Centrally purchased vaccines for the national immunisation programme for the NHS can only be ordered via ImmForm. Vaccines for use for the national immunisation programme are provided free of charge. National stock may also be used for catch-up vaccination of individuals of any age.</p> <p>Protocols for the ordering, storage and handling of vaccines should be followed to prevent vaccine wastage (see the Green Book Chapter 3).</p> <p>Also refer to All Wales Advisory document on Ordering Storage and Handling of Vaccines.</p>
Storage	<p>Store between +2°C to +8°C. Store in original packaging in order to protect from light. Do not freeze.</p> <p>Protocols for the storage and handling of vaccines should be followed to prevent vaccine wastage.</p> <p>ProQuad® must be administered immediately after reconstitution, to minimise loss of potency. Discard if the reconstituted vaccine is not used within 30 minutes.</p> <p>Priorix Tetra® must be administered promptly following reconstitution. Vaccines that are not used immediately</p>

	<p>must be kept in the refrigerator. If not used within 24 hours, the vaccine must be discarded.</p> <p>In the event of an inadvertent or unavoidable deviation of these conditions, vaccines that have been stored outside the conditions stated above should be quarantined and risk assessed for suitability of continued off-label use or appropriate disposal. Refer to All Wales Advisory document on Ordering Storage and Handling of Vaccines and Vaccine Incident Guidance.</p> <p>Contact the vaccine manufacturer where more specific advice is required about managing a temperature excursion.</p> <p>See 'MMP 427 Safe and Secure Management of Refrigerated Medicines and Vaccines SOP' for details of actions required in the event of a fridge temperature excursion.</p> <p>Any loss of vaccines due to expiry date or fridge failure/breaches in cold chain must be reported on ImmForm, to PTHB Immunisation co-ordinator (Powys.Immunisations@wales.nhs.uk), to the Medicines Management Team Info.MedicinesManagement.Powys@wales.nhs.uk and via PTHB Datix reporting system Once for Wales Reporting System.</p>
<p>Disposal</p>	<p>Follow local clinical waste policy and standard operating procedures to ensure safe and secure waste disposal.</p> <p>Equipment used for immunisation, including used vials, ampoules, or discharged vaccines in a syringe or applicator, should be disposed of safely in a UN-approved puncture-resistant sharps box, according to local waste disposal arrangements and NHS England guidance (HTM 07-01): safe and sustainable management of healthcare waste and guidance in the Welsh Health Technical Memorandum 07-01 Safer management of healthcare waste.</p>
<p>Drug interactions</p>	<p>The immunological response may be diminished in those receiving immunosuppressive treatment. Vaccination is recommended even if the antibody response may be limited (see criteria for exclusion).</p>

Reference Number: PGD 0250A
 Valid from: 02/02/2026
 Review Date: 31/05/2028
 Expiry Date: 30/11/2028

	<p>MMRV may be given at the same time or at any time before or after other inactivated vaccines. MMRV may attenuate the response to other live vaccines; see Table 11.4 (recommended time intervals when giving more than one live attenuated vaccine) in chapter 11 of the Green Book. Where protection against measles is required rapidly, other live vaccines should be given at any interval. The response may be suboptimal if yellow fever and MMR-containing vaccines are co-administered or given within a 4 week interval; an additional dose of MMR should be considered.</p> <p>There are no concerns from giving the MMRV vaccine to a child who has already been vaccinated with the monovalent varicella vaccine.</p> <p>If protection against measles is urgently required, then the benefit of protection from the vaccine outweighs the potential interference with a tuberculin (Mantoux) test. In this circumstance, the individual interpreting the negative tuberculin test should be made aware of the recent MMRV vaccination when considering how to manage that individual.</p> <p>When MMRV is given within 3 months of receiving blood products, such as immunoglobulin, the response to the measles component may be reduced. This is because such blood products may contain significant levels of measles-specific antibody, which could then prevent vaccine virus replication. Where possible, MMRV should be given at least one month before or deferred until 3 months after receipt of such products. If immediate measles protection is required in someone who has recently received a blood product, MMR vaccine should still be given. To confer longer-term protection, MMR should be repeated after 3 months.</p> <p>A detailed list of drug interactions associated with the MMRV vaccine is available from the product's SPC.</p>
<p>Identification and management of adverse reactions</p>	<p>Adverse reaction rates with ProQuad® are either similar to or less common after a second dose than after the first dose; incidence and severity of adverse reactions following a second dose with Priorix Tetra® are broadly similar.</p> <p>Adverse reactions are attributed to effective replication of the vaccine viruses, with subsequent mild illness.</p>

Reference Number: PGD 0250A
 Valid from: 02/02/2026
 Review Date: 31/05/2028
 Expiry Date: 30/11/2028

Events due to the measles component occur 6 to 11 days after vaccination. Events due to the mumps and rubella components usually occur 2 to 3 weeks after vaccination but may occur up to 6 weeks after vaccination. Adverse events due to the varicella component may occur up to a month following vaccination. Individuals with vaccine-associated symptoms are not infectious to others, although any varicella-type rash round the injection site should be covered. Transmission of varicella vaccine virus from immunocompetent vaccinees to immunocompromised close contacts has occasionally been documented, though the risk is low. Consider giving post-exposure prophylaxis to immunosuppressed contacts where the vaccinee has a disseminated rash (see the [varicella](#) chapter of the Green Book).

The most common adverse reactions are fever and injection site reactions including pain, swelling and erythema. Rash (including a measles-type and varicella-type rash) is also commonly reported.

Malaise, fever or a rash (or a combination of these) most commonly occur about a week after immunisation, lasting around 2 to 3 days.

Hypersensitivity reactions and anaphylaxis can occur but are very rare.

In case of an acute anaphylactic reaction occurring, adequate treatment provision must be available for immediate use: Anaphylaxis and resuscitation equipment including adrenaline (1 in 1000) injection and a working telephone.

In case of anaphylaxis:

- Refer to [adrenaline \(epinephrine\) PGD0017](#) and [anaphylaxis procedure](#)
- Request medical assistance urgently. If the GP is not immediately available dial 999 to transfer to A&E
- Ensure reaction is fully documented in patient notes
- Ensure all patient records are marked **ALLERGIC TO MMRV Vaccine**, stating the brand.
- The patient may be advised to wear a MedicAlert or similar device to alert other healthcare providers.
- Report via [Datix Once for Wales Reporting system](#)

Rare and more serious events

	<p>Febrile seizures, benign and short lived in nature are the most commonly reported neurological event following immunisation with MMRV and more commonly after the first compared to the second dose. Seizures occur between day 5 and 12 following vaccination in around 1 in 1000 children vaccinated with MMRV. The absolute risk of febrile seizures remains low.</p> <p>Encephalitis has been reported during post-marketing use of MMRV vaccines. In a few cases, fatal outcomes have been observed, especially in patients who were immunocompromised. Individuals, parents or carers should be instructed to seek prompt medical attention if they or their child experience symptoms suggestive of encephalitis such as loss or reduced levels of consciousness, convulsions or ataxia accompanied by fever and headache.</p> <p>Arthropathy (arthralgia or arthritis) has also been reported to occur rarely after MMR immunisation, probably due to the rubella component. If it is caused by the vaccine, it should occur between 14 and 21 days after immunisation. Where it occurs at other times, it is highly unlikely to have been caused by vaccination. The incidence rate is higher and the reaction more marked in adult females, though such reactions are generally well tolerated.</p> <p>ITP has occurred rarely following MMRV vaccination, usually within 6 weeks of the first dose and resolves spontaneously. The risk of developing ITP after MMRV vaccine is much less than the risk of developing it after infection with wild measles or rubella virus (see cautions).</p> <p>Further details on adverse reactions following MMRV vaccine can be found in the Green Book chapters on measles, mumps, rubella and varicella.</p> <p>A detailed list of adverse reactions associated with the MMRV vaccine is available from the product's SPC.</p>
<p>Reporting procedure of adverse reactions</p>	<p>Healthcare professionals and individuals, parents or carers are encouraged to report suspected adverse reactions to the Medicines and Healthcare products Regulatory Agency (MHRA) using the Yellow Card reporting scheme or by searching for MHRA Yellow Card in the Google Play or Apple App Store.</p>

Reference Number: PGD 0250A

Valid from: 02/02/2026

Review Date: 31/05/2028

Expiry Date: 30/11/2028

	<p>Any adverse reaction to a vaccine should be documented in the individual's record and the individual's GP should be informed.</p> <p>All significant adverse drug reactions and any administration errors must be recorded via the Once for Wales Reporting System.</p>
<p>Written information to be given to individual or parent (or carer)</p>	<p>Offer the marketing authorisation holder's patient information leaflet (PIL) provided with the vaccine.</p> <p>Priorix Tetra® PIL ProQuad® PIL</p> <p>For resources in accessible formats and alternative languages, please visit Find Public Health resources. Where applicable, inform the individual, parent or carer that large print, Braille or audio CD PILs may be available from emc accessibility by providing the medicine name and product code number, as listed on the product's SPC.</p> <p>Immunisation promotional material may be provided as appropriate:</p> <ul style="list-style-type: none"> • MMR for all • pre-school immunisations: guide to vaccinations (2 to 5 years) • measles: information for schools and healthcare centres • measles outbreak resources • Off-label vaccines - an introductory guide for healthcare professionals <p>NHS Wales: NHS 111 Wales - Vaccinations</p> <p>Further information for printing and website links suitable for the public can be found on the Public Health Wales intranet site Public Health Wales Immunisation and Vaccine Preventable Disease Programme, NHS 111 Wales and Health Information Resources.</p>
<p>Advice and follow up treatment</p>	<p>Inform the individual, parent or carer of possible side effects and their management.</p> <p>Advise about likely timing of and subsequent management of a fever.</p>

Reference Number: PGD 0250A

Valid from: 02/02/2026

Review Date: 31/05/2028

Expiry Date: 30/11/2028

	<p>Advise where relevant that pregnancy should be avoided for one month post vaccination.</p> <p>The individual, parent or carer should be advised to seek medical advice in the event of an adverse reaction and report this via the Yellow Card reporting scheme.</p> <p>When administration is postponed, advise the individual, parent or carer when to return for vaccination.</p> <p>Where applicable, advise the individual, parent or carer when the subsequent dose is due.</p> <p>Give appropriate advice if medication is used off-label.</p>
<p>Special considerations and additional information</p>	<p>Ensure there is immediate access to adrenaline (epinephrine) 1 in 1000 injection and access to a working telephone at the time of vaccination.</p> <p>All children with egg allergy should receive the MMRV vaccination as a routine procedure in primary care. Monovalent varicella vaccines and the varicella component of MMRV are not manufactured using eggs or egg-derived products. Although MMR-containing vaccines use egg-derived products in the manufacturing process, they are not used in the vaccine itself.</p> <p>ProQuad® contains porcine gelatine.</p> <p>Priorix® Tetra (GSK) does not contain porcine gelatine and can be offered as an alternative to ProQuad®. Health professionals should be aware of the need to order Priorix® Tetra when running clinics for relevant communities (see vaccines and porcine gelatine leaflet).</p> <p>MMRV vaccine is recommended when protection against measles, mumps, rubella or varicella (or a combination of the 4) is required. MMRV vaccine can be given irrespective of a history of measles, mumps, rubella or varicella infection or vaccination. There are no ill effects from vaccinating those who are already immune. If there is doubt about an individual's MMRV immune status, MMRV vaccine should still be given.</p>

Children with chronic conditions such as cystic fibrosis, congenital heart or kidney disease, failure to thrive or Down's syndrome are at particular risk from measles infection and should be immunised with an MMR-containing vaccine without delay.

Early vaccination due to travel, outbreak or contact with a probable or confirmed case of measles

The response to MMRV in infants is sub-optimal where the vaccine has been given before one year of age. Maternal antibodies may reduce the response to the first dose of vaccination up to the age of 18 months. To provide additional protection to those who fail to respond to the first dose, therefore, the second dose should not routinely be given below 18 months of age. A dose of MMRV given before the first birthday should be discounted. 2 further doses of MMR-containing vaccine should be given at the recommended ages in accordance with the routine schedule (see [dose and frequency of administration](#)).

Children over 12 months of age who require their second dose of MMRV to be given early due to travel, outbreak or contact with a case should have this dose brought forward to a minimum of 4 weeks after the first dose. If the child is aged under 18 months and a subsequent MMRV dose is given less than 3 months after the first MMRV dose, the dose should be repeated from 18 months of age and at least 4 weeks after the last dose, to ensure full protection.

Follow-up doses given after 18 months of age and at a minimum interval of 4 weeks from the previous MMRV dose(s) may be counted as valid.

Where applicable, the decision on when to vaccinate adults needs to take into consideration the past vaccination history, the likelihood of an individual remaining susceptible and the future risk of exposure and disease. See the Green Book chapters on [measles](#), [mumps](#), [rubella](#) and [varicella](#) for more information.

Children and adults coming from abroad may not have been immunised against measles, mumps, rubella or varicella. Unless there is a reliable history of appropriate immunisation, individuals should be assumed to be unimmunised. See [chapter 11](#) for more

information. Eligible individuals aged 18 months and over who have not received MMRV, or who received a dose of measles-containing vaccine before the age of one should receive 2 doses at least 4 weeks apart. An individual who has already received one dose of MMR from their first birthday should receive a second dose of MMR-containing vaccine (as MMR or MMRV as appropriate for their date of birth) to ensure that they are protected.

Individuals who have already received live (monovalent) varicella vaccine (including private market vaccination)

Individuals who have been previously immunised with 2 doses of the live varicella vaccine (see [varicella vaccine PGD0198](#)) should be considered to be fully protected against chickenpox. There is no requirement to check with the parent or carer for a history of varicella vaccination. MMRV may still be given to the child where the parent or carer expresses a wish for them to have the vaccine, even if 2 doses of monovalent varicella has been given (including under private market vaccination). However, no further additional protection is anticipated to be conferred from the additional MMRV doses.

Individuals who have previously received one dose of monovalent varicella vaccine require a further dose of varicella-containing vaccine in order to ensure full protection. This dose should be given as MMRV if the individual falls under the [inclusion criteria](#). Otherwise, complete the course with monovalent varicella vaccine – see the [varicella vaccine PGD0198](#).

Ensure an interval of at least 4 weeks is observed between live varicella vaccine and MMRV.

Chickenpox infection

There is no requirement to check for a history of chickenpox infection from the parent or carer of a child who responds to the catch-up offer. There are no safety concerns from giving MMRV to a child who has already had chickenpox infection.

Post exposure

Antibody responses to the rubella and mumps components of MMRV vaccine do not develop soon enough to provide effective prophylaxis after exposure to these infections. However, as vaccine-induced

	<p>measles antibody develops more rapidly than that following natural infection, MMRV vaccine should be used to protect susceptible contacts from suspected measles. To be effective against measles or varicella exposure, the vaccine must be administered very promptly and ideally within 3 days.</p> <p>Even where it is too late to provide effective post-exposure prophylaxis with MMRV, the vaccine can provide protection against future exposure. Therefore, contact with suspected measles, mumps, rubella or varicella provides a good opportunity to offer MMRV vaccine to previously unvaccinated, eligible individuals.</p> <p>If the individual is already incubating measles, mumps, rubella or varicella, MMRV vaccination will not exacerbate the symptoms. In these circumstances, individuals should be advised that a measles, mumps, rubella or varicella-like illness occurring shortly after vaccination is likely to be due to natural infection.</p> <p>Immunoglobulin may be indicated for contacts of measles who are infants, immunosuppressed or pregnant or for some varicella contacts. Provision of immunoglobulin is not covered by this PGD (see national measles guidelines or guidelines on post-exposure prophylaxis (PEP) for varicella or shingles respectively).</p> <p>NB Consult the PTHB Occupational Immunisation policy for staff requiring immunisation.</p>
<p>Records</p>	<p>Record consultation details as required by local procedures. In addition, the practitioner must ensure the following is recorded:</p> <ul style="list-style-type: none"> • that valid informed consent was given or a decision to vaccinate made in the individual’s best interests in accordance with the Mental Capacity Act 2005. Record name of representative who gave consent if appropriate • name of individual, address, date of birth and GP with whom the individual is registered (or record where an individual is not registered with a GP) • printed name and signature of immuniser (if recording on paper). If recording directly on WIS or another digital system, then name only as the signature will be replaced by electronic logging of the user.

	<ul style="list-style-type: none"> • medical and drug history taken, including any allergies and previous adverse events • name and brand of vaccine • date of administration • dose, form and route of administration of vaccine • quantity administered • batch number and expiry date • anatomical site of vaccination • any reasons for exclusion or referral • advice given, including advice given if the individual is excluded or declines immunisation • details of any adverse drug reactions and actions taken • administered via PGD (record PGD title and version number) <p>Records should be signed and dated (or password-controlled on e-records).</p> <p>All records should be clear, legible and contemporaneous.</p> <p>This information should be recorded in the individual's GP record. Where vaccine is administered outside the GP setting, appropriate health records should be kept and the individual's GP informed.</p> <p>When vaccine is administered to individuals under 19 years of age, the local Child Health Information Systems (CHIS) team (Child health records department) must be notified using the appropriate documentation or pathway as required by any local or contractual arrangement (based in Brecon hospital for under 5 years and Llandrindod hospital for school age).</p> <p>A record of all individuals receiving treatment under this PGD should also be kept for audit purposes in accordance with local policy.</p>
--	--

4. Key references

<p>Key references</p>	<p>MMRV vaccine</p> <ul style="list-style-type: none"> • Immunisation Against Infectious Disease: The Green Book measles, mumps, rubella and varicella, chapter 6 and chapter 11 www.gov.uk/government/collections/immunisation-against-infectious-disease-the-green-book • Introduction of a routine varicella (MMRV) vaccination programme (NHS system letter), published 31 October 2025 https://www.gov.uk/government/publications/introduction-of-a-routine-varicella-mmr-vaccination-programme • Summary of Product Characteristics for ProQuad[®], Merck Sharpe and Dohme, last updated 7 October 2025 https://www.medicines.org.uk/emc/product/101444/smpc • Summary of Product Characteristics for Priorix Tetra[®], GlaxoSmithKline, last updated 31 October 2025 https://www.medicines.org.uk/emc/product/101321/smpc • JCVI: Childhood varicella vaccination programme: JCVI advice, 14 November 2023 https://www.gov.uk/government/publications/childhood-varicella-vaccination-programme-jcvi-advice-14-november-2023 • MSD Medical Information, Personal communication (via email), 13 December 2023 • UKHSA national measles guidelines, last updated 25 July 2024. https://www.gov.uk/government/publications/national-measles-guidelines • Vaccination of individuals with uncertain or incomplete immunisation status, UKHSA
------------------------------	--

	<p>www.gov.uk/government/publications/vaccination-of-individuals-with-uncertain-or-incomplete-immunisation-status</p> <ul style="list-style-type: none"> • The National Society for Phenylketonuria (NSPKU) Medical Advisory Panel: Vaccines and PKU, issued 2 October 2024 https://nspku.org/download/vaccines-and-pku/ <p>General</p> <ul style="list-style-type: none"> • All Wales Advisory document on Ordering Storage and Handling of Vaccines 7th Edition September 2017 • NHSE Health Technical Memorandum 07-01: safe and sustainable management of healthcare waste, updated 7 March 2023 www.england.nhs.uk/publication/management-and-disposal-of-healthcare-waste-hm-07-01/ • National minimum standards and core curriculum for vaccination training www.gov.uk/government/publications/national-minimum-standards-and-core-curriculum-for-immunisation-training-for-registered-healthcare-practitioners • NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions, last updated March 2017 www.nice.org.uk/guidance/mpg2 • NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions, updated January 2018 www.nice.org.uk/guidance/mpg2/resources • UKHSA Immunisation Collection www.gov.uk/government/collections/immunisation • Vaccine Incident Guidance, last updated July 2022 https://www.gov.uk/government/publications/vaccine-incident-guidance-responding-to-vaccine-errors • Welsh Health Technical Memorandum 07-01 Safer management of healthcare waste
--	---

Appendix A: PGD indications for use for MMR, MMRV and monovalent varicella vaccine, in accordance with the individual’s age.

	6 to 9 months of age	9 months to less than 12 months of age	12 months of age and over
Monovalent varicella vaccine (V) PGD	Not recommended	Recommended vaccine for pre and post exposure to varicella	Recommended vaccine for pre and post exposure to varicella, where the individual is not eligible for the MMRV programme
MMR PGD	Indications for PGD unchanged: <ul style="list-style-type: none"> • early vaccination for travel to a measles endemic area • post exposure prophylaxis for measles • measles outbreaks 	Indications for PGD unchanged: <ul style="list-style-type: none"> • early vaccination for travel to a measles endemic area • post exposure prophylaxis for measles • measles outbreaks 	The individual is ineligible for the MMRV programme and either MMR protection is required in line with the vaccination of individuals with uncertain or incomplete immunisation algorithm or for travel, post exposure or outbreak
MMRV PGD	Not recommended	Alternative option for varicella pre and post-exposure, where Varivax® or Varilrix® is not available and protection is urgently required	<p>Routine vaccination at 12 and 18 months for children born on or after 1 January 2025; bringing MMRV status up to date for children born on or after 1 January 2020</p> <p>Individuals ineligible for MMRV, when MMR or monovalent varicella vaccine is not available* and who require urgent protection against MMR or V, such as in managing post-exposure varicella or measles outbreaks or administering an opportunistic catch-up dose of MMR vaccine.</p> <p>Where an individual requires both varicella vaccine and MMR vaccine at the same time, even if they are not eligible for MMRV in the routine programme.</p>

*MMR vaccine will be available for administration outside of the routine childhood programme (for example, for catching up older individuals, where date of birth is on or before 31 December 2019, who have not received 2 doses of MMR and are not eligible for MMRV). Therefore, after 1 January 2026, providers are expected to maintain stock of MMR vaccine for those ineligible for MMRV.

Reference Number: PGD 0250A
 Valid from: 02/02/2026
 Review Date: 31/05/2028
 Expiry Date: 30/11/2028

Appendix B – Staff Accredited to use the Patient Group Direction

Authorising Manager: I confirm that the practitioners named below have declared themselves suitably trained and competent to work under this PGD. I give authorisation on behalf of Powys Teaching Health Board or a Powys GP practice for the named healthcare professionals below who have signed the PGD to work under it. *The authorising manager must use the competency checklist (below).*

Practitioner: *By signing this PGD you are indicating that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each professional to practise only within the bounds of their own competence and professional code of conduct.*

I confirm that I have read and understood the content of this Patient Group Direction and that I am willing and competent to work to it within my professional code of conduct.

Printed name of health professional	Signature of health professional	Printed name of senior representative authorising health professional (Authorising Manager)	Signature of senior representative authorising health professional (Authorising Manager)	Date

The authorising manager should retain a copy of the list, which will be requested for audit purposes. This list should be kept by PTHB (or the provider organisation adopting an authorised version of the PGD) for 25 years after the PGD expires.

The healthcare professional should retain a copy of the document after signing.

Competency check list for manager or senior team lead to use as part of the authorising process for health professionals to work to a Patient Group Direction (PGD). Review of authorisation will take place on each PGD update and at the individual's annual PADR.

	Name: Role:	Sign / Initial	Further training identified (Y/N) Specify in " comments	Comments
1	The PGD sign off is for the following PGD:(document the exact title and PGD number) _____			
2	We have discussed the expiry of the PGD and are using a version accessed electronically			
3	The member of staff has the appropriate qualifications and professional registration as outlined in the PGD			
4	The Patient Group Direction has been read in full by the staff member			
5	The identified training has been completed as specified in the PGD and is in date			
6	We have discussed some examples of inclusion criteria and exclusion criteria			
7	The staff member is confident in the administration method and doses			

Staff member print & sign name		Date
Manager or senior team lead to print & sign name		Date

Please send a copy of this completed form to individual's line manager and to the staff member, in conjunction with the PGD Appendix B authorisation sheet. A copy of this form should also be kept by service lead in the training folder

